



[Previous abstract](#) | [Graphical version](#) | [Text version](#) | [Next abstract](#)

Session LC25 - Tunneling in High-Tc Superconductors I.

ORAL session, Tuesday afternoon, March 23

Room 171W, GWCC

[LC25.03] Quasiparticle Tunneling and Andreev Reflection Study of the Pseudogap in Underdoped $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ and $\text{YBa}_2\text{Cu}_3\text{O}_{6+x}$

J. Y.T. Wei, C.-T. Chen, C.-C. Fu, N.-C. Yeh (Department of Physics, California Institute of Technology, Pasadena, CA 91125), R.P. Vasquez (Center for Space Microelectronics Technology, Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109)

Scanning tunneling and point-contact spectroscopy measurements are performed on epitaxial-film and single-crystal samples of underdoped $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ and $\text{YBa}_2\text{Cu}_3\text{O}_{6+x}$ to study the pseudogap characteristics as a function of temperature. The measurements are made between 4.2K and 100K, to understand how the size, symmetry, and character of the superconducting order parameter evolve through the critical temperature T_c and below the pseudogap temperature T^* . In particular, the $2\Delta_0/k_B T_c$ and $2\Delta^*/k_B T^*$ ratios are determined, where Δ_0 and Δ^* are the respective gap-maxima. The occurrence of a zero-bias peak in the tunneling conductance is interpreted as a signature of d-wave pairing symmetry, while an Andreev-reflection behavior in the point-contact spectra is understood as a direct manifestation of the particle-particle nature of the pseudogap. The results are discussed in terms of recent theoretical models, including pair-density fluctuations and Bose-Einstein condensation of pre-formed pairs.

■ Part I of program listing

superconduction